PCT.

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

(11) International Publication Number:

WO 99/20237

A61K 7/16

A1

(43) International Publication Date:

29 April 1999 (29.04.99)

(21) International Application Number:

PCT/IB97/01634

(22) International Filing Date:

17 October 1997 (17,10,97)

(71) Applicant (for all designated States except US): ZAKRYTOE AKTSIONERNOE OBSCHESTVO OSTIM [RU/RU]; per. Sechenovsky, 6-3, Moscow, 119034 (RU).

(72) Inventors: and

(75) Inventors/Applicants (for US only): RUDIN, Vsevolod Nikolaevich [RU/RU]; ul. Petrozavodskaya, 15–2–199, Moscow, 125502 (RU). ZUEV, Vladislav Petrovich [RU/RU]; ul. Tallinskaya, 2–316, Moscow, 123426 (RU). KOMAROV, Vladimir Fedorovich [RU/RU]; Shipilovsky proezd, 53–5–606, Moscow, 115582 (RU). MELIKHOV, Igor Vitallevich [RU/RU]; ul. Krupskol, 19–55, Moscow, 117331 (RU). MINAEV, Vladimir Vasillevich [RU/RU]; Naberezhnaya Tarasa Shevchenko, 5–471, Moscow, 121248 (RU). ORLOV, Andrei Yurlevich [RU/RU]; ul. Kantimirovskaya, 12–1–471, Moscow, 121357 (RU). MISHIN, Anatoly Aleksandrovich [RU/RU]; ul. Obrucheva, 3–2–77, Moscow, 117421 (RU). BOZHEVOLNOV, Viktor Evgenievich [RU/RU]; ul. Bolshaya Kommunisticheskaya, 24/28–45, Moscow, 109004 (RU).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: STOMATIC COMPOSITION

(57) Abstract

This invention relates to the field of medicine, and in particular to the stomatology and may be used for preventive treatment and combatting caries, parodenitis and paradentosis. The stomatic composition on the basis of hydroxyapatite also comprises abrasive materials, humectants, thickeners, surfactants, flavouring agents, and a number of optional ingredients. The composition comprises hydroxyapatite in the form of particles of ultra finely divided hydroxyapatite with an average size by length (1), width (d) and thickness (h) of about 1 x d x h = 0.06 μ m x 0.015 μ m; x 0.005 μ m is capable to stimulate reparative osteogenesis processes and possessing a high bioactivity and specific pharmacological activity. The ultra finely divided hydroxyapatite is present in the composition in the amount of 0.1 % to 50 % by weight. The ultra finely divided hydroxyapatite is a synthetic hydroxyapatite which contains about 99.9 % of Ca₁₀(PO₄)6(OH)₂ by weight. The stomatic composition can be used to cure microdefects of the basic substance of the dental enamel, can be applied to prepare toothpastes, toothcreams and gels and can also be included as a component of chewing gum, pastilles, tooth elixir, formulations to rinse mouth.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

ES FI FR GA GB Herzegovina GE GH	Gabon United Kingdom	LS LT LU LV MC	Lesotho Lithuania Luxembourg Latvia	SI SK SN SZ	Slovenia Slovakia Senegal
FR GA GB Herzegovina GE	France Gabon United Kingdom	LU LV	Luxembourg Latvia	SN	Slovakia Senegal
GA GB Herzegovina GE	Gabon United Kingdom	LV	Latvia		Senegal
GB Herzegovina GE	United Kingdom			S7.	
Herzegovina GE	6	MC			Swaziland
	Geomie		Monaco	TD	Chad
GH	Georgia	MD	Republic of Moldova	TG	Togo
	Ghana	MG	Madagascar	T.)	Tajikistan
GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
so GR	Greece		Republic of Macedonia	TR	Turkey
HU	Hungary	ML	Mali	TT	Trinidad and Tobago
IE	Ireland	MN	Mongolia	UA	Ukraine
IL	Israel	MR	Mauritania	UG	Uganda
· IS	Iceland	MW	Malawi	US	United States of America
. IT	Italy	MX	Mexico	UZ	Uzbekistan
ican Republic JP	Japan	NE	Niger	VN	Viet Nam
KE	Kenya	NL	Netherlands	YU	Yugoslavia
KG		NO	Norway	zw	Zimbabwe
re KP	Democratic People's	NZ	New Zcaland	2	Zimbab we
	Republic of Korea				
KR					
KZ					•
ıblic LC	•				
LK					·
.!	Li	KR Republic of Korea KZ Kazakstan blic LC Saint Lucia LI Liechtenstein LK Sri Lanka	KR Republic of Korea PT KZ Kazakstan RO blic LC Saint Lucia RU LI Liechtenstein SD LK Sri Lanka SE	KR Republic of Korea PT Portugal KZ Kazakstan RO Romania blic LC Saint Lucia RU Russian Federation LI Liechtenstein SD Sudan LK Sri Lanka SE Sweden	KR Republic of Korea PT Portugal KZ Kazakstan RO Romania blic LC Saint Lucia RU Russian Federation LI Liechtenstein SD Sudan LK Sri Lanka SE Sweden

WO 99/20237 PCT/IB97/01634

STOMATIC COMPOSITION

Description

Field of the Invention

This invention relates to the field of medicine, and in particular to the field of stomatology and may be used for preventive treatment and curing of caries, parodenitis and paradentosis.

Prior art

For these above-captioned purposes, stomatic compositions comprising hydroxyapatite (HA) have found an extensive application in the stomatologic practice.

There are certain compositions having a favourable effect including synthetic HA containing 92 to 97% $Ca_{10}(PO_4)_6(OH)_2$, 3 to 6% H_2O and 0,3% $CaCO_3$ with an average particle size of 1 to 15 μm .

Such a stomatic composition, for instance, according to Patent EP 0344832 cl. A61K 7/16, comprises save the stated HA, water-soluble casein material or sodium trimetaphosphate, as an anti-caries agent and also other well_known_ingredients_which_depend_upon_the_forms_of_the_product_manufactured, such as various humectants, binding thickeners, surfactants, flavouring agents.

The known stomatic composition (EP 03442746 cl. A61K 7/18, publ.23.11.89) supplementary includes a fluorine-containing compound in the form of NaP or sodium monophluorphosphate as an anti-caries agent.

The amount of HA present in the stomatic composition is in the range of 1 to 50%, usually 2 to 20% by weight of the stomatic composition. The stomatic composition comprises some other ingredients: humectants, thickeners, surfactants

and flavouring agents commonly known to those skilled in the art in all formulations of such products.

However, the stomatic compositions stated possess a relatively poor anti-caries effect and is not useful in the preventive medicine and in the treatment of inflammatory-destructive diseases of paradentium tissues.

Disclosure of the invention

It is an object of the present invention to create a stomatic composition comprising compounds capable to cure microdefects of the basic substance of the dental enamel to combat caries developing (e.g. to provide an anti-caries activity) and to prevent the spread of such inflammable-destructive diseases of paradentium tissues as paradenitis and paradentosis, and also compounds capable to stimulate reparative osteogenesis processes and possessing high bioactivity and specific pharmacological activity.

It is a further object of the invention to create a stomatic composition being identic to the basic substance of the dental enamel in its substance contents and crystalline parameters, as the acid formed in the materials covering dental surfaces causes destruction of mineral hydroxyapatyte out of which enamel is composed and has a result due to which calcium ion loss occurs.

The task surprisingly has been solved in a composition as defined in claim 1.

A preferred composition having a more pronounced effect in view of the improvements obtained according to the invention comprises particles of hydroxyapatite with an average particle size in length (1), width(d) and thickness(h) of about 1 = 0,06 μ m +/- 50 %, d = 0,015 μ m +/- 50 % and h = 0,005 μ m +/- 50 %.

A most preferred composition having a surprisingly superior effect in view of the improvements obtained according to the invention comprises particles of hydroxyapatite with an average particle size in length (1).

width(d) and thickness(h) of l about 0,06 $\mu m,$ d about 0,015 $\mu m,$ h about 0,005 $\mu m.$

Being introduced into the composition, HA possesses osteo-reparative properties and contains preferably about $100\% Ca_{10} (PO_4)_6 (OH)_2$.

The specific surface of HA used in the composite advantageously is 100 to 150 $\ensuremath{\text{m}}^{2}/g\,.$

The amount of HA present in the oral composition of the present invention is in the range of 0,1% to 50%, preferably from 0,1% to 25%, and most preferably from about 0,2% to 20% by weight of the oral composition.

'The composition reacts to a change in the biochemical environment, for instance a rapid dissolvement of ultra finely divided HA occurs when the pH is decreasing, that provides an active utilization of Ca and PO, - ions in the osteogenesis process: the size and configuration of the inventive crystals are adapted to the maximum to the dental enamel structure, which is mostly composed of HA, that suggests its use in the osteo-reparative process as a building material.

The ultra finely divided HA possesses a high adhesive-sorption activity to the dental enamel and to microdefects on its surface, that favour the preventive measures preventing the spread of caries disease and also possesses a high sorption activity in respect to proteins and aminoacids of paradentium tissues, that stimulates an active preventive treatment of the inflammable-destructive diseases such as paradenitis and paradentosis.

Moreover, the stomatic composition of the present invention will contain other conventional ingredients in addition to HA possessing osteo-reparative properties, whose introduction into the composition depends on the form of the product. For instance, in the case of an oral product in the form of dentifrice paste, cream or gel, the product will comprise a liquid phase containing humectants and binding thickeners which act to maintain the particulate solid

abrasive and HA crystals in the form of stable suspension in the liquid phase.

Surfactants and flavouring agents are also usual ingredients for various inventive embodiments of oral compositions.

The humectants usually used are glycerol or sorbitol. However, other humectants may be used according to the invention including polyethyleneglycol, propyleneglycol, lactitol and hydrogenated corn syrop. The amount of humctant will generally range from about 0% to 85% by weight of product. The remainder of the liquid phase will consist substantially of water. The liquid phase can be water or a non-aqueous composition.

As binding agents and thickeners, various substances can be used such as sodium carboxymethylcellulose, sodium hydroxyethylcellulose and xanthan gum. Natural gum bindings can be included such as gum tragacanth, gum karaya of Irish moss, etc. Any mixture of binding agents and thickeners can be also used. The amount of bindings and thickeners usually included into the oral composition is in the range of 0% to 10% by weight of the oral composition.

Moreover, any materials as widely disclosed in the literature generally also might be used for the invention as surfactants, i.e. surfactants like sodium lauryl sulphate, dodecylbenzene sulphonate and sodium lauryl sarcosinate. Other anionic surfactants also can be used as well as cationic and amphoteric and non-ionic surfactants. Surfactants are generally present in the composition in the amount of 0% to 5% by weight of the oral composition.

Flavours that are generally used in the oral compositions are those based on oils of spearmint and peppermint and might be used for the invention. Examples of other flavouring materials used are menthol, clove, wintergreen, eucalyptus and aniseed. A preferable amount of flavours is from 0% to 5% by weight in respect to the oral composition.

As abrasive materials, silica dioxide of various modifications, aluminium oxide, calcium carbonate, dicalcium phosphate anhydrite, dicalcium phosphate dihydrate, sodium metaphosphate insoluble in water, and thereof mixtures may be used. The amount of abrasive materials ranges from 0.0% to 25%. The oral composition may include a wide variety of optional ingredients. These include antimicrobial and antiplaque agents for example chlorhexidine or 2,4,4-trichloro-2hydroxy-diphenyl ether, or zink compounds (see EPA-161898) anti-tartar ingredients such as condensed phosphates, e.g. alkali et al pyrophosphates, hexametaphosphatesor polyphosphates (see US-A-4 515772 and US-A-4 627977) or zink citrates (see US-A-4 100269), sweetening agents such as saccharin. Preservatives such as formalin, sodium benzoate. colouring agents (for instance titanium dioxide) or pHcontrolling agents, such as acid base or buffer agents the oral composition may also include agents enhancing the gingivitis system of the mouth cavity and representing extracts of various natural plants such as urtica, millefolium, chamomilla hypericum, salvia, etc. in the aqueous or aqueous-alcoholic forms.

The stomatic composition depending on its form (dentifrice paste, cream or gel) is maintained in contact with the tissue of the oral cavity from 15 sec to 12 hours.

The following examples of dentifrice pastes and gel comprising synthetic ultra finely divided HA possessing osteo-reparative properties as described above illustrate the invention. Percentages and parts of the components are by weight.

Belowstanding preferred embodiments of the invention are shown in its composition.

Examples N1 and 2.

Toothpaste prepared from the following ingredients.

$\mathcal{C}_{i} = \{ \mathbf{c}_{i} \in \mathcal{C}_{i} \mid i \in \mathcal{C}_{i} \mid i \in \mathcal{C}_{i} \} $		
Ingredi	ents,%	
Example	1	2
V	•	
Ultra finely divided		
Hydroxyapatite	0,2	2,0
Silica aerogel	22,0	15,0
Sodium carboxymethylcellulose	1,0	1,0
Glycerol distilled	20,0	20,0
Sorbitol	20,0	17,0
Titanium dioxide	0,6	0,5
Sodium benzoate	0,4	0,6
Aqueous-alcohol extract	•	
of chamomilla	1,0	0,8
Aqueous-alcohol extract	:	
of hypericum	1,0	0,8
Sodium saccharin	0,1	0,06
Flavour	1,0	1,3
Sodium lauryl sulphate	1,5	1,5
Water	to 100,0	to 100,0

Examples N 3 to 7 $\begin{tabular}{ll} \begin{tabular}{ll} \hline \begin{tabular}{ll} \be$

	Ingredie	nts,%	:		
Example	3	· 4	5	6	7
	· .		` :		
Ultra finely divided				:	
hydroxyapatite		2,5		2,5	2,5
Silica aerogel Sodium	17,0	17,0	17,0	17,0	17,0
hydroxyethylcellulose Sodium	1,6	÷	-	1,6	-
carboxymethylcellulose		1,1	1,1	· ·	0,9
Sorbitol	20,0	20,0	16,0	20,0	•
Glycerol distilled	20,0	20,0	20,0	20,0	20,0
Polyethyleneglycol	-	-	5,0	-	-
Sodium lauryl sulphate	1,5	1,5	1,5	1,5	1,5
Tetrasodium pyrophospha	te -	1,5	-	-	-
Tetrapotassium	•	•			
pyrophosphate	-	-	-	2,5	_
Sodium trimetaphosphate	-	-	2,0	.	-
Zinc citrate trihydrate	<u>-</u>	-	-	-	0,5
Titanium dioxide	0,6	0,6	0,6	0,6	0,6
Sodium benzoate	0,5	0,5	0,6	=	-
Formalin	-	-	<u>-</u>	0,05	0,05
Aqueous-alcohol extract					
of salvia	0,5	0,5	-	-	-
Aqueous-alcohol extract					
of millefolium	0,9	0,9	0,5	0,5	-
Aqueous-alcohol extract					
of chamomilla	-	-	1,0	0,7	· -
Triclosan	-	-	0,2	-	0,2
Sodium saccharin	0,06	0,06	0,06	0,06	0,06

R

Flavour	1,0	1,	0	1,0	1,0	1,0
Water			in	all exa	mple to	100,0

Examples N8 and 9
Gel preventing paradenitis.

'ı Ing	gredients,%	8			
Example	8	9			
Ultra finely divided	•		•		
hydroxyapatite	5,0	4,0			
Sodium hydroxyethylcellulose	2,0	2,5			
Silica aero	5,0	-			
Glycerol distilled	10,0	-			
Sorbitol	25,0	45,0			
Sodium benzoate	0,5	-			
Triclosan	- .	0,3			
Flavour	0,2	0,15			
Sodium lauryl sulphate	0,2	0,15			
Sodium saccharin	0,07	0,07			
Water	to 100,00	to 100,00			

Industrial application

The stomatic composition can be used to cure microdefects of the basic substance of the dental enamel, e.g. to prevent the spread of caries, and is also useful for

preventive measures avoiding the spread of inflammabledestructive diseases of paradentium tissues, such as pardenitis and paradentosis.

The stomatic composition can be used in the form of tooth pastes, tooth creams and gels. Moreover, the composition can be included as a component in jewing gum; pastilles, tooth elixir and formulations to rinse mouth.

The stomatic composition according to the invention is capable to stimulate reparative osteogenesis processes and possessing a high bioactivity and specific pharmacological activity. Moreover, this composition is useful for combatting dental caries and to prevent the spread of such inflammable-destructive diseases of paradentium tissues as paradenitis and paradentosis, on the basis of hydroxyapatite also optionally comprising abrasive materials, humectants, thickeners, surfactants, flavouring agents, and a number of optional ingredients.

Claims:

0,0001 µm.

- 1. A stomatic composition characterised in that it comprises particles of hydroxyapatite with an average particle size in length (1), width (d) and thickness (h) of: 1 from about 0,2 μ m to about 0,01 μ m, d from about 0,1 μ m to about 0.001, and h from about 0.1 μ m to about
- 2. The stomatic composition according to claim 1 characterised in that it comprises particles of hydroxyapatite with an average particle size in length (1), width(d) and thickness(h) of about $l=0.06~\mu m$ +/- 50 %, $d=0.015~\mu m$ +/- 50 % and $h=0.005~\mu m$ +/- 50 %.
- 3. The stomatic composition according to claim 1 characterised in that it comprises particles of hydroxyapatite with an average particle size in length (1), width(d) and thickness(h) of about 1 = 0,06 μ m, d = 0,015 μ m, h = 0,005 μ m.

4. A stomatic composition

characterised in

that it comprises particles of hydroxyapatite having a specific surface of hydroxiapatite from about 100 m $^2/g$ to about 150 m $^2/g$.

5. The stomatic composition according to one of claims 1 to 4 characterized in that it comprises said hydroxyapatite particles ultra finely divided.

- 6. The composition according to one of claims 1 to 5 characterised in that the ultra finely divided hydroxyapatite particles are present in the composition in an amount of 0,1% to 50% by weight.
- 7. The composition according to one of claims 1 to 6 characterised in that the ultra finely, divided hydroxyapatite is a synthetic hydroxyapatite which contains 99,9% of $Ca_{10}(PO_4)_6(OH)_2$ by weight.
- 8. The composition according to one of claims 1 to 7 further characterised by at least one substance of the group consisting of
 - humectants in a range from about 0% to 85% by weight,
 - bindings and thickeners a range of 0% to 10% by weight,
 - abrasive materials in a range from 0.0% to 25%,
 Surfactants in a range from 0% to 5% by weight,
 - Flavours in a range from 0% to 5% by weight.
- 9. The composition according to one of claims 1 to 8 further characterised by agents enhancing the gingivitis system of the mouth cavity and comprising extracts of natural plants including at least one of the group consisting of urtica, millefolium, chamomilla hypericum, salvia, etc. in the aqueous an in the aqueous-alcoholic form.
- 10. The composition according to one of claims 1 to 9 further characterised by antimicrobial and anti-plaque agents.

INTERNATIONAL SEARCH REPORT

I national Application No

		· i	101718 31702001
A. CLASS IPC 6	SIFICATION OF SUBJECT MATTER A61K7/16		*
According	to International Patent Classification(IPC) or to both national classi	fication and IPC	
_	S SEARCHED	Realion dire ii C	
	ocumentation searched (classification system followed by classific	ation symbols)	
IPC 6	A61K		
Documenta	ation searched other than minimum documentation to the extent tha	t such documents are includ	led in the fields searched
			:
Electronic	data base consulted during the international search (name of data	base and, where practical, s	earch terms used)
		<u> </u>	
	ENTS CONSIDERED TO BE RELEVANT		
Category ²	Citation of document, with indication, where appropriate, of the r	elevant passages	Relevant to claim No.
X	EP 0 786 245 A (SANGI CO., LTD) 1997	30 July	1-10
	see the whole document	•	
х	EP 0 664 133 A (AKTIONERNOE OBS ZAKRYTOGO TIPA "OSTIM") 26 July see claim 1; example 1		1
E	WO 98 18719 A (AKTSIONERNOE OBSO ZAKRYTOGO TIPA "OSTIM") 7 May 19 see claims 1,14		1
	·		
		•	
-			
-		••	
			· .
		· · · · · · · · · · · · · · · · · · ·	
Funth	ner documents are listed in the continuation of box C.	X Patent family me	mbers are listed in annex.
° Special car	tegories of cited documents :		hed after the international filing date
"A" docume conside	ant defining the general state of the art which is not ered to be of particular relevance	cited to understand t	not in conflict with the application but the principle or theory underlying the
	locument but published on or after the international		r relevance; the claimed invention
"L" docume	are that which may throw doubts on priority claim(s) or scied to establish the publication date of another	involve an inventive	d novel or cannot be considered to step when the document is taken alone
citation	or other special reason (as specified) and referring to an oral disclosure, use, exhibition or	cannot be considere	r relevance; the claimed invention d to involve an inventive step when the ad with one or more other such docu-
other n	neans nt published prior to the international filing date but	ments, such combine in the art.	ation being obvious to a person skilled
	an the priority date claimed	"8" document member of Date of mailing of the	international search report
٠.	July 1998	16/07/199	·
Name and m	nailing address of the ISA	Authorized officer	
	European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk		
**	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Glikman,	J-F

INTERNATIONAL SEARCH REPORT

Information on patent family members

rational Application No PCT/IB 97/01634

Patent document cited in search repor	t	Publication date		Patent family member(s)	Publication date
ÈP 786245	A	30-07-1997	JP	9202717 A	05-08-1997
EP 664133	Α	26-07-1995	RU WO	2077329 C 9503074 A	20-04-1997 02-02-1995
WO 9818719	Α	07-05-1998	NONE		

Form PCT/ISA/210 (patent family annex) (July 1992)